

The tool-use network supports actions independent of the acting body part

Florencia Martinez Addiego^{1*+}, Yuqi Liu^{1,2*}, Kyungji Moon^{1,3}, Elizabeth Shytle¹, Lénia Amaral^{1,4}, Caroline O'Brien¹, Sriparna Sen^{1,5}, Maximilian Riesenhuber^{1,6}, Jody C. Culham⁷, and Ella Striem-Amit¹

* These authors contributed equally to this work
+ Contact email: fam53@georgetown.edu

¹Department of Neuroscience, Georgetown University Medical Center, Washington, DC 20057, USA; ²Institute of Neuroscience, Key Laboratory of Primate Neurobiology, CAS Center for Excellence in Brain Sciences and Intelligence Technology, Chinese Academy of Sciences, Shanghai, China; ³Department of Neurology, Georgetown University Medical Center, Washington, DC 20057, USA; ⁴Faculty of Psychology and Educational Sciences, University of Coimbra, Portugal; ⁵Department of Psychology, University of Pittsburgh, Pittsburgh, PA, USA; ⁶Center for Neuroengineering, Georgetown University, Washington, DC 20057, USA; ⁷Department of Psychology, University of Western Ontario, London, Ontario, N6A 5C2, Canada



Organization of actions in the brain


- The sensorimotor system is broadly organized **somatotopically**.
- Yet, we can execute the same action with **different body parts**, suggesting there is a **motor plan** that **generalizes across certain action parameters**.
- Indeed, higher-level action regions (e.g. the **parietal lobe** and the **premotor dorsal areas**) represent more **abstract action information across motor parameters**.
- These association action areas support reaching and grasping actions when performed by other body parts. Does this generalization extend to **non-evolutionarily based actions**, like **tool use**?
- We tested if the tool-use network supports the generalization of action-type across body parts (hand and foot).

MOTIVATING QUESTION:

Does the brain support generalization across acting body parts for tool-use actions?

Methods and Approach


fMRI experiment of action execution:
n = 22 right-handed controls (5 excluded)
n = 8 individuals born without hands (IDs; 2 excluded)




The tool-use network generalizes across body parts

Hand tool-use selective regions

Hand tool-use regions show consistent tool-use preference in foot actions



Hand tool-use areas have a shared neural code between the hand and the foot




Hand tool-use areas generalize across body parts in motor association areas

Conclusions

- Tool-use areas** defined from the hand execution data show a **preference for tool-use in foot actions, regardless of sensorimotor experience**
- The PMd, SPL (area2 and 7pc), and LOTC have a **shared neural code across the hand and the foot** in typically-developed participants
- Temporal analyses reveal a **hierarchical dissociation** between **lower-level M1** which groups by body part and **higher-level association areas** which instead tend to cluster by action-type

Temporal dynamics patterns cluster by action in association areas



LOOKING FOR A
POSTDOC

FUNDING

Edwin H. Richard and Elisabeth Richard von Matsch
Distinguished Professorship in Neurological Diseases (to E.S.A)